

University of Groningen

Reciprocal associations between symptom levels of disturbed grief, posttraumatic stress, and depression following traumatic loss

Lenferink, Lonneke I. M.; Nickerson, Angela; de Keijser, Jos; Smid, Geert E.; Boelen, Paul A.

Published in:
Clinical Psychological Science

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lenferink, L. I. M., Nickerson, A., de Keijser, J., Smid, G. E., & Boelen, P. A. (Accepted/In press). Reciprocal associations between symptom levels of disturbed grief, posttraumatic stress, and depression following traumatic loss: A four-wave cross-lagged study. *Clinical Psychological Science*.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

**Reciprocal associations between symptom levels of disturbed grief, posttraumatic stress,
and depression following traumatic loss: A four-wave cross-lagged study**

Running title: Reciprocal links between grief-related symptoms

Lonneke I.M. Lenferink^{1,2}, Angela Nickerson³, Jos de Keijser¹, Geert E. Smid^{4,5,6} and Paul
A. Boelen^{2,4,5}

¹ Department of Clinical Psychology and Experimental Psychopathology, Faculty of Behavioral and Social Sciences, University of Groningen, Grote Kruisstraat 2/1, 9712 TS Groningen, the Netherlands. E-mail L.I.M. Lenferink: l.i.m.lenferink@rug.nl; E-mail J. de Keijser: a.de.keijser@rug.nl

² Department of Clinical Psychology, Faculty of Social Sciences, Utrecht University, P.O. Box 80140, 3508 TC, Utrecht, the Netherlands. E-mail P.A. Boelen: p.a.boelen@uu.nl

³ School of Psychology, University of New South Wales, Sydney, Australia. E-mail A. Nickerson: a.nickerson@unsw.edu.au

⁴ Foundation Centrum '45, Nienoord 5, 1112 XE Diemen, the Netherlands. E-mail G.E. Smid: g.smid@centrum45.nl

⁵ Arq Psychotrauma Expert Group, Nienoord 5, 1112 XE, Diemen, the Netherlands

⁶ University of Humanistic Studies, Kromme Nieuwegracht 29, 3512 HD Utrecht, the Netherlands.

Corresponding author: Lonneke I. M. Lenferink, Department of Clinical Psychology and Experimental Psychopathology, Faculty of Behavioral and Social Sciences, University of

Groningen, Grote Kruisstraat 2/1, 9712 TS Groningen, the Netherlands. E-mail:

l.i.m.lenferink@rug.nl, telephone: (+31) (0) 50-3639189.

Conflict of interest

All authors declare that there is no conflict of interest.

Funding

The Victim Fund subsidized this work.

Contributions

LL and PB designed the study. LL collected the data. LL and AN conducted the statistical analyses. LL and PB wrote the draft of the manuscript. All authors critically reviewed and improved draft versions of the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgements

We would like to thank Eline Piersma, MSc., and Daniëlle Trip, MSc., for their support with data-collection. We would also like to thank Victim Support the Netherlands for their support with recruiting participants.

In press: Clinical Psychological Science, accepted for publication on April 18, 2019

Abstract

Objective

Disturbed grief, operationalized as persistent complex bereavement disorder (PCBD), correlates with, yet differs from posttraumatic stress disorder (PTSD) and depression symptoms. However, knowledge about temporal associations between these symptoms is limited. We aimed to enhance our understanding of the aetiology of loss-related distress by examining temporal associations between PCBD, PTSD, and depression symptom levels.

Methods

Dutch people (N=172) who lost significant other(s) in a plane disaster completed questionnaires for PCBD, PTSD, and depression 11, 22, 31, and 42 months post-disaster.

Results

Cross-lagged analyses revealed that changes in PCBD symptom levels have a greater impact on changes in symptom levels of PTSD and depression than vice versa.

Conclusion

Our findings contradict the notion that PTSD and depression symptoms should be addressed before grief in treatment. Pending replication of our findings in clinical samples, we tentatively conclude that screening and treatment of grief symptoms has potential value in preventing long-lasting distress.

Keywords: persistent complex bereavement disorder, grief, depression, posttraumatic stress, disaster, trauma

Persistent complex bereavement disorder (PCBD) is included as condition for further study in the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5; American Psychiatric Association, 2013). PCBD symptoms encompass grief reactions, including preoccupation with the deceased and feeling numb since the loss, that persist for at least one year following the death of a significant other and cause significant impairments in functioning. A meta-analysis suggests that 9.8% of people confronted with the non-violent loss (e.g., illness) of a significant other is at risk for developing clinically relevant grief reactions (henceforth referred to as PCBD; Lundorff, Holmgren, Zachariae, Farver-Vestergaard, & O'Connor, 2017). Experiencing the death of someone significant that happens unexpectedly and/or has a violent cause heightens the risk of PCBD and comorbid posttraumatic stress disorder (PTSD) and depression (Boelen, de Keijser, & Smid, 2015; Currier, Holland, & Neimeyer, 2006; Kaltman & Bonanno, 2003). This combination of symptoms has also been referred to as traumatic grief (Smid et al., 2015).

PCBD shares symptoms with PTSD (e.g., re-experiencing, avoidance symptoms (Maccallum & Bryant, 2018; Maercker & Znoj, 2010; Nickerson et al., 2014; O'Connor, Lasgaard, Shevlin, & Guldin, 2010)) and depression (e.g., sad mood, emptiness, fatigue, and suicidal ideation (Boelen, van den Bout, & de Keijser, 2003; Boelen, Reijntjes, Djelantik, & Smid, 2016; Djelantik, Smid, Kleber, & Boelen, 2017; Maccallum, Malgaroli, & Bonanno, 2017; Shear et al., 2011)). Re-experiencing symptoms in PCBD differ from those in PTSD such that for PCBD re-experiencing comprises both positive and negative memories, whereas for PTSD re-experiencing of negative memories is key (Maercker & Znoj, 2010).

Furthermore, avoidance in PCBD serves to avert painful thoughts and feelings related to the irreversibility of the loss, whereas avoidance in PTSD is used to prevent thoughts and memories related to the traumatic event and the recurrence of threat or danger. Depression often coincides with inactivity, whereas yearning in PCBD seems to be associated with

passivity as well as activity, such as searching behaviour (Stroebe & Schut, 1999). In addition, a sad mood in depression is related to low self-esteem, whereas a sad mood in PCBD is related to longing for the deceased (Shear et al., 2011).

Although research has shown that some disturbed grief symptoms resemble symptoms of PTSD and depression, studies employing for instance factor or latent class analysis have shown that these syndromes are correlated, but distinguishable (Boelen, van de Schoot, van den Hout, de Keijser, & van den Bout, 2010; Bonanno et al., 2007; Lenferink, de Keijser, Smid, Djelantik, & Boelen, 2017; Li & Prigerson, 2016; Nickerson et al., 2014; O'Connor et al., 2010). However, knowledge about the temporal associations between these symptoms is limited. Gaining more insight in these associations is important for our understanding of the aetiology of post-loss distress and could inform us about symptoms that need to be dealt with first in treatment of comorbid loss-related symptoms. To our knowledge, only two studies have examined associations between PCBD and PTSD symptom levels over time. One study did so using four waves of data and lower level mediation analysis in elderly spousal bereaved people (O'Connor, Nickerson, Aderka, & Bryant, 2015). Another study examined these associations in a community sample of bereaved people using two waves of data and cross-lagged panel analysis (Djelantik, Smid, Kleber, & Boelen, 2018). Findings from both studies indicated that changes in PCBD symptom levels have a greater impact on changes in PTSD symptom levels than vice versa. These findings are important in attesting to the predictive validity of PCBD and run counter to prevailing notions that in treatment of combined symptoms of grief and traumatic distress, traumatic distress needs to be targeted before disturbed grief reactions should be addressed (Lindy, Green, Grace, & Titchener, 1983; Raphael & Martinek, 1997; Sprang, 2001). However, it should be noted that both prior studies, examining reciprocal relationships, relied on non-clinical bereaved samples (Djelantik

et al., 2018; O'Connor et al., 2015). Therefore, caution is warranted in generalizing these findings to clinical samples.

While these findings are important, it is yet unclear to what extent they generalize to people who have lost a significant other due to a violent cause. This is important given research findings suggesting that people exposed to a violent loss are over 5 times more likely to develop comorbid PCBD and PTSD reactions (vs. resilient response) than people exposed to a non-violent loss (Djelantik et al., 2017). The temporal relationship between PCBD and depression has also not been explored in prior research; consequently, it is still unclear whether increases in PCBD symptom levels precede worsening of depression symptom levels or whether elevated depression symptom levels predicts an exacerbation of grief reactions. Accordingly, we examined the reciprocal associations between symptom levels of PCBD, PTSD, and depression in a non-clinical sample of people who suffered one or multiple significant losses in a plane disaster in 2014.

Aims of the study

We are the first to investigate the temporal relationship between self-reported PCBD and PTSD symptom levels in people exposed to a violent loss of a significant other. We expanded prior work by also exploring the reciprocal associations between PCBD and depression and the reciprocal associations between PCBD, PTSD, *and* depression symptom levels. In line with previous findings (Djelantik et al., 2018; O'Connor et al., 2015), we expected that changes in PCBD symptom levels would predict changes in PTSD symptom levels but not vice versa. Because no prior studies have examined the temporal associations between PCBD and depression symptom levels, no specific hypotheses were formulated regarding these associations over time.

Material and Methods

Participants

The MH17 disaster took place on July 17 2014. In total 298 people were on board when the airplane was shot down in Ukrainian airspace. All people, including 193 Dutch people, died. Dutch adult people who had lost one or multiple family member(s) and/or friend(s) in this disaster were invited to take part in this longitudinal online survey study. Four waves of data are included in this study. One prior study has been published based on these four waves of data, in which we employed latent class growth modelling to examine individual differences in longitudinal symptom-profiles (Lenferink, Nickerson, de Keijser, Smid, & Boelen, 2018). People were recruited using a variety of strategies, including invitations sent by peer-support and governmental support organisations, media attention, and snowball sampling (for details see our earlier work (Lenferink, de Keijser, Smid, Djelantik, & Boelen, 2017).

In total 172 people completed at least one wave and were included in the analyses. The majority of the participants were women ($N = 102$, 59.3%) and about two-third ($N = 117$, 68.4%) had at least a university degree. The participants' mean age, at the time the disaster took place, was 51.55 ($SD = 15.51$) years. Fifty-four people (31.8%) lost one significant other. Most people lost multiple significant others; 58 people (34.1%) lost two, 22 people (12.9%) lost three, 32 (18.8%) lost four, 2 people (1.2%) lost five, and 2 people (1.2%) lost six significant others. Regarding relationship to the deceased, for people who suffered multiple losses, the most intimate relationship to the deceased was reported here, ordered from child, through spouse, through parent, through sibling, through other not previously specified significant other (cf., Lenferink et al., 2017). Forty-eight people (28.4%) lost at least a child, 2 people (1.2%) at least a spouse, 15 people (8.9%) at least a parent, 48 people (28.4%) at least a sibling, 30 people (17.8%) at least another family member, 12 people (7.1%) at least a friend, and 14 people (8.3%) other than spouse, family member, or a friend (e.g., neighbour or colleague). The mean PCBD, PTSD, and depression scores fell within the

subclinical range (see Table 1; for more details about clinical cut-offs see Lenferink et al., 2018).

Procedures

The first assessment (referred to as Wave 1 (W1)) took place from May 2015 through January 2016. The second assessment (W2) took place from April 2016 through December 2016. The third assessment (W3) took place from February 2017 through May 2017. The fourth assessment (W4) took place from January 2018 through March 2018. The mean time since loss was 10.73 (SD = 1.77) months for W1, 21.91 (SD = 1.47) months for W2, 31.31 (SD = 0.98) months for W3, and 41.66 (SD = 0.59) months for W4. In total, 66 participants (38.4%) completed all four waves. A local ethical board approved this study. All participants provided written informed consent.

Measures

PCBD symptoms as defined in DSM-5 (American Psychiatric Association, 2013) were assessed with 16 items of the 18-item Traumatic Grief Inventory-Self Report (Boelen & Smid, 2017). The frequency of each symptom (e.g., “I had trouble accepting the loss”) during the previous month was rated on scales ranging from 1 (“never”) through 5 (“always”). In the instruction we referred to ‘the loss of your loved one(s) due to the plane crash’ as anchor event. Total score ranged from 16 – 80. The TGI-SR has adequate psychometric properties (Boelen & Smid, 2017; Boelen, Djelantik, de Keijser, Lenferink, & Smid, 2018). Cronbach’s alphas in the current study were $>.88$ for each wave.

PTSD symptoms as per DSM-5 were assessed¹ with the 20-item PTSD Checklist for DSM-5 (Blevins, Weathers, Davis, Witte, & Domino, 2015; Boeschoten, Bakker, Jongedijk,

¹ We used the PCL-5 to assess PTSD symptoms at all waves, except W1. Twenty-five randomly chosen participants filled in another PTSD measure at W1 that was used by collaborators in this project. We excluded these data and treated these as missing completely at random in the analyses.

& Olf, 2014). Participants rated to what extent they experienced each symptom (e.g., “In the past month, how much were you bothered by: “Trouble remembering important parts of the stressful experience?””) during the preceding month, on scales ranging from 0 (“not at all”) through 4 (“extremely”). In the instruction we referred to ‘the loss of your loved one(s) due to the plane crash’ as anchor event. The PCL-5 has adequate psychometric properties (Blevins et al., 2015; Krüger-Gottschalk et al., 2017). Cronbach’s alphas in the current study were $>.91$ for each wave.

Symptoms of depression were assessed with the 16-item Quick Inventory of Depressive Symptomatology-Self Report (Rush et al., 2003). The frequency of each symptom (e.g., “Feeling irritable”) during the preceding 7 days was rated on scales with four answer options ranging from 0 through 3². As recommended by Rush et al. (Rush et al., 2003), the nine items were summed to yield a total depression score. Psychometric properties of the QIDS-SR are adequate (Rush et al., 2003). Cronbach’s alphas in the current study were $>.75$ for each wave.

Statistical analyses

Three series of cross-lagged panel models were conducted, using Mplus version 8 (Muthén & Muthén, 1998-2017), to test the reciprocal relationships between symptom levels of PCBD, PTSD, and/or depression (represented by summed scores on the TGI-SR, PCL-5, and QIDS-SR, respectively) across the four waves. In the first round of analyses, we included only PCBD and PTSD data, the second round focused on PCBD and depression data only, and the third round included PCBD, PTSD, *and* depression data. Each cross-lagged model consisted of autoregressive paths to take into account symptom persistence over time (e.g., the associations between PCBD symptom levels at W1, W2, W3, and W4). Based on previous

² For item 12 (i.e., “thoughts of deaths or suicide”), we removed answer option 3 for ethical reasons.

research (Boelen et al., 2010; Bonanno et al., 2007), we also took into account associations between PCBD, PTSD, and depression symptom levels at each wave, by allowing these distress measures to correlate in the model (e.g., PCBD at W1 with PTSD at W1 in the first round of analyses). Cross-lagged effects were estimated to test the predictive effect of a distress measure at one wave (e.g., symptom levels of PCBD at W1) to the other distress measure at a subsequent time point (e.g., symptom levels of PTSD at W2), while controlling for the autoregressive effects and associations between the distress measures at each wave.

In accord with recommendations put forth by Little (2013), the most parsimonious cross-lagged model was preferred, i.e., a model with reduced set of estimated parameters that fits the data well. We therefore compared the statistical fit of the unconstrained model (Model 1) with increasingly constrained models (i.e., assuming associations are equal) in consecutive steps. In Model 2, we constrained the autoregressive paths to be equal for the distress measures separately. In Model 3, we constrained the autoregressive paths plus the cross-lagged effects of PCBD on PTSD and depression and of PTSD and depression on PCBD, and in the last round of analyses of PTSD on depression and of depression on PTSD, to be equal. In Model 4, we also constrained the associations between the distress measures at the same wave to be equal across the waves. Selection of the most parsimonious model was based on the following fit statistics: 1) the root mean square error of approximation (RMSEA) value of $<.08$ representing acceptable fit and value of $<.05$ excellent fit, and the comparative fit index (CFI) and Tucker-Lewis Index (TLI) with values $>.90$ indicating acceptable fit. In a final round of analyses, we added covariates to the final models simultaneously by regressing the baseline symptom levels on these covariates. Based on a systematic review and meta-analysis from Heeke et al. (2019), we included gender (0 = man, 1 = woman), kinship to the deceased (0 = other than child/spouse, 1 = child/spouse), and educational level (0 = lower than university, 1 = university) as covariates. Missing data on symptom levels of PCBD, PTSD,

and depression were handled using full information maximum likelihood estimation. All significance tests were two-tailed ($\alpha = .05$).

Results

Preliminary analyses

Participants completing W1 but not completing W2, W3, or W4 did not significantly differ from those completing these waves in terms of background characteristics, except that those not completing W2, W3, or W4 were more distantly related to the deceased than those completing these waves. There were also no significant differences in symptom levels of PCBD and PTSD. However, participants not completing W3 and W4 had lower depression levels than those completing these waves (see our prior work for more details Lenferink et al. (2018)).

Table 1 shows means, SDs, and score ranges of, and correlations between symptom levels at all waves. PCBD, PTSD, and depression symptoms were significantly and positively correlated at each time point.

Table 1

Means, standard deviations, and correlations of PCBD, PTSD, and depression symptom levels at W1 to W4

		N	Mean	SD	Range	PCBD			PTSD				Depression			
						W2	W3	W4	W1	W2	W3	W4	W1	W2	W3	W4
PCBD	W1	167	40.73	10.36	21-78	.69	.71	.57	.81	.65	.67	.59	.64	.57	.47	.40
	W2	93	38.14	12.03	16-77		.82	.70	.61	.88	.76	.74	.48	.74	.58	.56
	W3	100	35.80	12.59	16-79			.82	.70	.76	.86	.78	.57	.66	.72	.71
	W4	102	35.16	11.25	16-77				.54	.63	.67	.79	.49	.59	.51	.62
PTSD	W1	139	19.17	14.06	0-77					.74	.81	.69	.78	.68	.61	.59
	W2	93	18.35	15.00	0-77						.82	.80	.60	.83	.65	.65
	W3	102	15.87	14.60	0-80							.86	.67	.77	.77	.76
	W4	103	16.26	12.98	0-74								.58	.77	.71	.80
Depression	W1	164	7.70	4.83	0-25									.74	.66	.62
	W2	95	7.58	4.89	0-22										.76	.75
	W3	101	7.17	5.06	0-22											.85
	W4	102	6.55	4.29	0-20											

Note. All correlations are significant at $p < .001$. PCBD = persistent complex bereavement disorder; PTSD = posttraumatic stress disorder.

Cross-lagged models

The fit indices of each cross-lagged model are shown in Supplemental Material Table 1. The models in which we constrained both the autoregressive and cross-lagged effects to be equal (Model 3) yielded the best fit, as indicated by the lowest RMSEA values and acceptable CFI and TLI values, and were therefore selected as our final models.

Temporal associations among symptom levels of PCBD, PTSD, and depression

Figures 1, 2, and 3 show the standardized autoregressive and cross-lagged effects of the final models. In Supplemental Material Table 2, the unstandardized final model results are shown. PCBD symptom levels significantly predicted PTSD symptom levels at a subsequent time point, but not vice versa (see Figure 1). PCBD symptom levels significantly predicted depression symptom levels at a subsequent time point, but not vice versa (see Figure 2). Figure 3 shows that PCBD and depression symptom levels significantly predicted PTSD symptom levels at a subsequent time point. PTSD symptom levels significantly predicted depression symptom levels at a subsequent time point. PTSD and depression symptom levels did not significantly predict PCBD symptom levels at a subsequent time point. In an additional round of analyses, we included the covariates gender, educational level, and kinship to the deceased into the cross-lagged models. This did not lead to meaningful differences in the results, i.e., the same significant cross-lagged effects were found³.

³ For the model including PCBD and PTSD symptom levels, the standardized significant ($p < .05$) cross-lagged effects for PCBD predicting subsequent PTSD varied from 0.14 through 0.18. For the model including PCBD and depression symptom levels, the standardized cross-lagged effects for PCBD predicting depression varied from 0.11 through 0.14. For the model including all symptom levels, standardized cross-lagged effects for PCBD predicting subsequent PTSD varied from 0.12 through 0.15, for depression predicting subsequent PTSD the effects varied from 0.15 through 0.17, and for PTSD predicting subsequent depression the effects varied from 0.18 through 0.21.




Figure 1. PCBD and PTSD symptoms over time: standardized autoregressive and cross-lagged effects

Note. Correlations between symptoms at same wave are not shown. Dashed lines represent non-significant associations. PCBD = persistent complex bereavement disorder; PTSD = posttraumatic stress disorder. * $p < .05$ ** $p < .01$ *** $p < .001$




Figure 2. PCBD and depression symptoms over time: standardized autoregressive and cross-lagged effects

Note. Correlations between symptoms at same wave are not shown. Dashed lines represent non-significant associations. PCBD = persistent complex bereavement disorder. * $p < .05$ ** $p < .01$ *** $p < .001$




Figure 3. PCBD, PTSD, and depression symptoms over time: standardized cross-lagged effects

Note. Autoregressive effects and correlations between symptoms at same wave are not shown. Dashed lines represent non-significant associations. PCBD = persistent complex bereavement disorder; PTSD = posttraumatic stress disorder. * $p < .05$ ** $p < .01$ *** $p < .001$

Discussion

In this study, we examined temporal associations between symptom levels of PCBD, PTSD, and depression in people that experienced the loss of significant others due to a plane disaster, using four waves of self-report questionnaire data. A first main finding was that symptom levels of PCBD predicted PTSD symptom levels whereas, conversely, PTSD symptom levels did not predict PCBD symptom levels. This finding accords with results from two prior recent studies (Djelantik et al., 2018; O'Connor et al., 2015). While these prior studies relied on people who mostly experienced a non-violent death, used different time intervals between the measurement occasions, and used different instruments to tap PCBD and PTSD (Djelantik et al., 2018; O'Connor et al., 2015), the consistency between our findings and these prior studies supports the robustness of these findings. Importantly, although our significant cross-lagged effects were relatively small, our findings run counter to frequently cited claims that symptoms of traumatic distress that may emerge following loss block the alleviation of symptoms of grief (Lindy et al., 1983; Raphael & Martinek, 1997; Sprang, 2001). However, our study, as well as prior studies examining reciprocal associations between psychopathology symptom levels (Djelantik et al., 2018; O'Connor et al., 2015), were not exclusively focused on bereaved people with clinically relevant psychopathology-levels. Pending replication of the findings in clinical samples, caution should be applied in generalizing our findings to clinical samples.

Expanding prior work, ours was the first to study temporal associations between PCBD and depression symptom levels. These associations mirrored the outcomes of the analyses with symptoms of PCBD and PTSD, such that results indicated that PCBD symptom levels significantly predicted depression symptom levels later in time, whereas depression symptom levels did not predict subsequent PCBD symptom levels. This finding complements treatment studies indicating that grief-specific treatment effectively reduce PCBD and

depression symptom levels, while depression-based treatments, such as interpersonal psychotherapy or use of antidepressants, are less effective in reducing PCBD symptom levels (Reynolds et al., 1999; Shear et al., 2014). Furthermore, the finding that PCBD symptoms predicted depression symptoms contradicts claims that alleviation of depressive symptoms enables bereaved people to grieve more effectively (Zisook, Shuchter, Pedrelli, Sable, & Deaciuc, 2001).

Notably, we also found that PCBD symptoms no longer predicted depression symptoms at a subsequent time point, once PTSD symptoms were taken into account in the last round of analyses. In general, prior research has also shown that when multiple outcomes are taken into account, this may yield a greater understanding of psychological responses after bereavement than studies examining a limited number of possible responses (Infurna & Luthar, 2017; Lenferink et al., 2018). More specifically, this result suggests that PTSD rather than PCBD symptoms predict depression symptoms over time. Yet, we also found that PCBD and depression symptoms predicted PTSD symptoms, while PTSD symptoms also predicted depression symptoms at consecutive waves of assessment. The bidirectional associations between PTSD and depression symptoms over time has also been found in people exposed to stressful events other than traumatic losses (Cheng, Liang, Fu, & Liu, 2018; Horesh, Lowe, Galea, Uddin, & Koenen, 2015). Consequently, our findings may indicate that an increase in PCBD symptoms predicts worsening of an intertwined cluster of PTSD and MDD symptoms over time. Our study yielded no evidence that PTSD and depression symptom levels are reciprocally related to PCBD symptom levels as these change in the first three to four years following the traumatic loss of significant others. These findings offer first evidence suggesting that it might be relevant to focus more on PCBD, rather than PTSD and depression, in screening and treating psychopathology among people heading toward the first anniversary of the unnatural death of a loved one. When our findings are replicated in clinical

samples, that would suggest that identification and treatment of elevated PCBD severity in this period may help to prevent the exacerbation of PTSD and depression levels at later periods in the bereavement process.

For each round of analyses we found that constraining the autoregressive and cross-lagged effects to be equal across the waves yielded the best statistical fit. This indicates that, over time, the predictive value of prior symptom levels remains quite stable on average. In other words, it seems unlikely that the strength of the temporal associations between PCBD, PTSD, and depression change in the first four years after the traumatic loss of a significant other.

The finding that increased PCBD symptoms seem to precede increases in PTSD and depression symptoms (and not vice versa, as has been argued previously (Lindy et al., 1983; Raphael & Martinek, 1997; Sprang, 2001; Zisook et al., 2001)) might be explained from at least two theoretical perspectives. Firstly, from the perspective of stress sensitization theory it could be argued that over time people become more sensitized to stress after confrontation with a life stressor. In other words, levels of stress needed to trigger onset of affective symptoms drastically decrease after each (Monroe & Harkness, 2005; Post, 2016; Stroud, 2018). From this perspective, bereaved people with elevated PCBD levels might be more prone to experience a worsening of PTSD and depression symptoms after confrontation with loss-related stimuli (Smid et al., 2015). For instance, the plane disaster that caused the death of their loved one(s) in the current sample has repeatedly received media-attention, which, in turn, could have led to elevated distress-levels (Neria et al., 2007; van der Velden, van der Meulen, Lenferink, & Ijzermans, 2018). Exposure to these stressors might be particularly harmful for those with elevated PCBD symptoms. Secondly, from the perspective of cognitive-behavioural models of disturbed grief (Boelen, van den Hout, & van den Bout, 2006), it could be argued that persistent yearning for the lost person's presence and

difficulties accepting his/her absence keep the person away from processing the most distressing and traumatizing features of the circumstances of the loss and, consequently, maintain PTSD symptoms (e.g., intrusive memories) resulting from the insufficiently processed memories of the circumstances (Boelen, 2010, 2017). Integration of information about the loss, including the relationship with the deceased, and the loss-related thoughts, memories, and feelings, with existing autobiographical knowledge has been argued to be critical for recovery from loss from a cognitive-behavioural perspective (Boelen et al., 2006). From this same cognitive-behavioural perspective, avoidance of social and occupational activities out of the belief that these are not fulfilling anymore since the loss (i.e., depressive avoidance) might play a role (Boelen & van den Bout, 2010). That is, over time social withdrawal may fuel depressive symptoms (Hong, Hasche, & Bowland, 2009). Consequently, those who have more difficulty in re-engaging in meaningful social activities after the plane disaster, might be more at risk to develop elevated depression symptoms.

Several limitations need to be kept in mind while interpreting our findings. In the current study we focused on PCBD levels according to the DSM-5 (American Psychiatric Association, 2013). Findings from recent studies suggest that PCBD per DSM-5 and prolonged grief disorder (PGD) as defined in the 11th edition of the International Classification of Diseases (WHO, 2018) perform differently when it comes to identification of clinical cases as well as concurrent and prospective validity (Boelen, Lenferink, Nickerson, & Smid, 2018; Maciejewski, Maercker, Boelen, & Prigerson, 2016; Mauro et al., 2017). Therefore, our findings do not necessarily generalize to disturbed grief as represented by PGD ICD-11 criteria. Furthermore, the majority of our sample was exposed to multiple losses simultaneously caused by a man-made disaster, most people were highly educated, and reported subclinical psychopathology-levels. This limits the generalizability to people confronted with other types of loss, lower educational level, and with clinical levels of

psychopathology. Lastly, we used self-report questionnaire data, instead of clinical diagnostic interviews. As a result, we were unable to formally diagnose PCBD, PTSD, and depression. Furthermore, because we used self-report questionnaire data the reported symptom levels might be overestimated (cf., Engelhard et al., 2007).

Notwithstanding these limitations, our findings show that changes in PCBD levels in people exposed to a traumatic loss three to four years earlier precede change in PTSD and depression levels and not vice versa. This runs counter to the widely held notion in the literature on grief treatment that PTSD and depression symptoms interfere with (and, thus, precede) grief reactions and that, among people getting stuck in recovery from loss, these should be addressed before targeting the grief. When our findings are replicated in clinical bereaved samples, our findings point to the value of screening and treatment of PCBD symptoms after the violent loss of a significant other in preventing long-lasting comorbid symptoms of PCBD, PTSD, and depression.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. Arlington, VA: American Psychiatric Publishing.
- Blevins, C. A., Weathers, F. W., Davis, M. T., Witte, T. K., & Domino, J. L. (2015). The posttraumatic stress disorder checklist for DSM-5 (PCL-5): Development and initial psychometric evaluation. *Journal of Traumatic Stress, 28*(6), 489-498.
- Boelen, P. A., van den Bout, J., & de Keijser, J. (2003). Traumatic grief as a disorder distinct from bereavement-related depression and anxiety: A replication study with bereaved mental health care patients. *American Journal of Psychiatry, 160*, 1339-1341.
- Boelen, P. A. (2017). "It feels as if she might return one day": A sense of unrealness as a predictor of bereavement-related emotional distress. *Estudios De Psicología, 38*(3), 734-751.
- Boelen, P. A., Lenferink, L. I. M., Nickerson, A., & Smid, G. E. (2018). Evaluation of the factor structure, prevalence, and validity of disturbed grief in DSM-5 and ICD-11. *Journal of Affective Disorders, 240*, 79-87. doi:10.1016/j.jad.2018.07.041
- Boelen, P. A., Reijntjes, A., Djelantik, A. A. A. M. J., & Smid, G. E. (2016). Prolonged grief and depression after unnatural loss: Latent class analyses and cognitive correlates. *Psychiatry Research, 240*, 358-363.
- Boelen, P. A., & Smid, G. E. (2017). The traumatic grief inventory self report version (TGI-SR): Introduction and preliminary psychometric evaluation. *Journal of Loss and Trauma, 22*(3), 196-212.

- Boelen, P. A., van de Schoot, R., van den Hout, M. A., de Keijser, J., & van den Bout, J. (2010). Prolonged grief disorder, depression, and posttraumatic stress disorder are distinguishable syndromes. *Journal of Affective Disorders, 125*(1-3), 374-378.
- Boelen, P. A., van den Hout, M. A., & van den Bout, J. (2006). A cognitive-behavioral conceptualization of complicated grief. *Clinical Psychology: Science and Practice, 13*(2), 109-128.
- Boelen, P. A., & van den Bout, J. (2010). Anxious and depressive avoidance and symptoms of prolonged grief, depression, and posttraumatic stress-disorder. *Psychologica Belgica, 50*(1), 49.
- Boelen, P. A., Djelantik, A. A. A. M. J., de Keijser, J., Lenferink, L. I. M., & Smid, G. E. (2018). Further validation of the traumatic grief inventory-self report (TGI-SR): A measure of persistent complex bereavement disorder and prolonged grief disorder. *Death Studies, 1-14*. doi:10.1080/07481187.2018.1480546
- Boelen, P. A. (2010). A sense of 'unrealness' about the death of a loved-one: An exploratory study of its role in emotional complications among bereaved individuals. *Applied Cognitive Psychology, 24*(2), 238-251. doi:10.1002/acp.1557
- Boelen, P. A., de Keijser, J., & Smid, G. (2015). Cognitive-behavioral variables mediate the impact of violent loss on post-loss psychopathology. *Psychological Trauma: Theory, Research, Practice, and Policy, 7*(4), 382-390.
- Boeschoten, M. A., Bakker, A., Jongedijk, R. A., & Olf, M. (2014). *PTSS checklist voor de DSM-5 (PCL-5)*. [Dutch PTSD checklist for DSM-5] Unpublished manuscript.

- Bonanno, G. A., Neria, Y., Mancini, A., Coifman, K. G., Litz, B., & Insel, B. (2007). Is there more to complicated grief than depression and posttraumatic stress disorder? A test of incremental validity. *Journal of Abnormal Psychology, 116*(3), 342-351.
- Cheng, J., Liang, Y., Fu, L., & Liu, Z. (2018). Posttraumatic stress and depressive symptoms in children after the Wenchuan earthquake. *European Journal of Psychotraumatology, 9*(1), 1472992. doi:10.1080/20008198.2018.1472992
- Currier, J. M., Holland, J. M., & Neimeyer, R. A. (2006). Sense-making, grief, and the experience of violent loss: Toward a mediational model. *Death Studies, 30*(5), 403-28.
- Djelantik, A. A. A. M. J., Smid, G. E., Kleber, R. J., & Boelen, P. A. (2017). Symptoms of prolonged grief, post-traumatic stress, and depression after loss in a Dutch community sample: A latent class analysis. *Psychiatry Research, 247*, 276-281.
- Djelantik, A. A. A. M. J., Smid, G. E., Kleber, R. J., & Boelen, P. A. (2018). Do prolonged grief disorder symptoms predict post-traumatic stress disorder symptoms following bereavement? A cross-lagged analysis. *Comprehensive Psychiatry, 80*, 65-71.
- Engelhard, I. M., van den Hout, M. A., Weerts, J., Arntz, A., Hox, J. J., & McNally, R. J. (2007). Deployment-related stress and trauma in Dutch soldiers returning from Iraq. Prospective study. *British Journal Psychiatry, 191*(2), 140-145.
- Heeke, C., Kampisiou, C., Niemeyer, H., & Knaevelsrud, C. (2019). A systematic review and meta-analysis of correlates of prolonged grief disorder in adults exposed to violent loss. *European Journal of Psychotraumatology*, Doi: 10.1080/20008198.2019.1583524

- Hong, S., Hasche, L., & Bowland, S. (2009). Structural relationships between social activities and longitudinal trajectories of depression among older adults. *The Gerontologist, 49*(1), 1-11.
- Horesh, D., Lowe, S. R., Galea, S., Uddin, M., & Koenen, K. C. (2015). Gender differences in the long-term associations between posttraumatic stress disorder and depression symptoms: Findings from the Detroit neighborhood health study. *Depression and Anxiety, 32*(1), 38-48. doi:10.1002/da.22267
- Infurna, F. J., & Luthar, S. S. (2017). The multidimensional nature of resilience to spousal loss. *Journal of Personality and Social Psychology, 112*(6), 926-947.
doi:10.1037/pspp0000095
- Kaltman, S., & Bonanno, G. A. (2003). Trauma and bereavement: Examining the impact of sudden and violent deaths. *Journal of Anxiety Disorders, 17*(2), 131-147.
- Krüger-Gottschalk, A., Knaevelsrud, C., Rau, H., Dyer, A., Schäfer, I., Schellong, J., & Ehring, T. (2017). The German version of the posttraumatic stress disorder checklist for DSM-5 (PCL-5): Psychometric properties and diagnostic utility. *BMC Psychiatry, 17*(1), 379.
- Lenferink, L. I. M., de Keijser, J., Smid, G. E., Djelantik, A. A. A. M. J., & Boelen, P. A. (2017). Prolonged grief, depression, and posttraumatic stress in disaster-bereaved individuals: Latent class analysis. *European Journal of Psychotraumatology, 8*(1)
doi:10.1080/20008198.2017.1298311

- Lenferink, L. I. M., Nickerson, A., de Keijser, J., Smid, G. E., & Boelen, P. A. (2018). Trajectories of grief, depression, and posttraumatic stress in disaster-bereaved people. *Depression and Anxiety*, doi:10.1002/da.22850
- Li, J., & Prigerson, H. G. (2016). Assessment and associated features of prolonged grief disorder among Chinese bereaved individuals. *Comprehensive Psychiatry*, 66, 9-16.
- Lindy, J. D., Green, B. L., Grace, M., & Titchener, J. (1983). Psychotherapy with survivors of the Beverly Hills supper club fire. *American Journal of Psychotherapy*, 37(4), 593-610.
- Little, T. D. (2013). *Longitudinal structural equation modeling*. New York: The Guilford Press. Retrieved from <https://rug.on.worldcat.org/oclc/815382938>
- Lundorff, M., Holmgren, H., Zachariae, R., Farver-Vestergaard, I., & O'Connor, M. (2017). Prevalence of prolonged grief disorder in adult bereavement: A systematic review and meta-analysis. *Journal of Affective Disorders*, 212(786-794), 138-149.
- Maccallum F, & Bryant, R. A. (2018). Symptoms of prolonged grief and posttraumatic stress following loss: A latent class analysis. *The Australian and New Zealand Journal of Psychiatry*, , 4867418768429. doi:10.1177/0004867418768429
- Maccallum, F., Malgaroli, M., & Bonanno, G. A. (2017). Networks of loss: Relationships among symptoms of prolonged grief following spousal and parental loss. *Journal of Abnormal Psychology*, 126(5), 652-662.
- Maciejewski, P. K., Maercker, A., Boelen, P. A., & Prigerson, H. G. (2016). "Prolonged grief disorder" and "persistent complex bereavement disorder", but not "complicated grief", are one and the same diagnostic entity: An analysis of data from the Yale bereavement study. *World Psychiatry*, 15(3), 266-275.

- Maercker, A., & Znoj, H. (2010). The younger sibling of PTSD: Similarities and differences between complicated grief and posttraumatic stress disorder. *European Journal of Psychotraumatology, 1*, 5558. doi:10.3402/ejpt.v1i0.5558
- Mauro, C., Shear, M. K., Reynolds, C. F., Simon, N. M., Zisook, S., Skritskaya, N., . . . Glickman, K. (2017). Performance characteristics and clinical utility of diagnostic criteria proposals in bereaved treatment-seeking patients. *Psychological Medicine, 47*(4), 608-615. doi:10.1017/S0033291716002749
- Monroe, S. M., & Harkness, K. L. (2005). Life stress, the “kindling” hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychological Review, 112*, 417-445.
- Muthén, L. K., & Muthén, B. O. (1998-2017). *Mplus user's guide. eighth edition*. Los Angeles: Muthén & Muthén.
- Neria, Y., Gross, R., Litz, B., Maguen, S., Insel, B., Seirmarco, G., . . . Marshall, R. D. (2007). Prevalence and psychological correlates of complicated grief among bereaved adults 2.5–3.5 years after September 11th attacks. *Journal of Traumatic Stress, 20*(3), 251-262.
- Nickerson, A., Liddell, B. J., Maccallum, F., Steel, Z., Silove, D., & Bryant, R. A. (2014). Posttraumatic stress disorder and prolonged grief in refugees exposed to trauma and loss. *BMC Psychiatry, 14*(1), 106.
- O'Connor, M., Nickerson A., Aderka I. M., & Bryant R. A. (2015). The temporal relationship between change in symptoms of prolonged grief and posttraumatic stress following old age spousal bereavement. *Depression and Anxiety*, doi:10.1002/da.22349

- O'Connor, M., Lasgaard, M., Shevlin, M., & Guldin, M. B. (2010). A confirmatory factor analysis of combined models of the Harvard Trauma Questionnaire and the Inventory of Complicated Grief-Revised: Are we measuring complicated grief or posttraumatic stress? *Journal of Anxiety Disorders, 24*(7), 672-9.
- Post, R. M. (2016). Epigenetic basis of sensitization to stress, affective episodes, and stimulants: Implications for illness progression and prevention. *Bipolar Disorders, 18*(4), 315-324. doi:10.1111/bdi.12401
- Raphael, B., & Martinek, N. (1997). Assessing traumatic bereavement and posttraumatic stress disorder. In J. Wilson, & T. Keane (Eds.), *Assessing psychological trauma and PTSD* (pp. 373-395). New York: Guilford Press.
- Reynolds, C. F. 3rd., Miller, M. D., Pasternak, R. E., Frank, E., Perel, J. M., Cornes, C., . . . Kupfer, D. J. (1999). Treatment of bereavement-related major depressive episodes in later life: A controlled study of acute and continuation treatment with nortriptyline and interpersonal psychotherapy. *American Journal of Psychiatry, 156*, 202-208.
- Rush, A. J., Trivedi, M. H., Ibrahim, H. M., Carmody, T. J., Arnow, B., Klein, D. N., . . . Keller, M. B. (2003). The 16-item quick inventory of depressive symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): A psychometric evaluation in patients with chronic major depression. *Biological Psychiatry, 54*(5), 573-583.
- Shear, M. K., Wang, Y., Skritskaya, N., Duan, N., Mauro, C., & Ghesquiere, A. (2014). Treatment of complicated grief in elderly persons. A randomized clinical trial. *JAMA Psychiatry, 71*(11), 1287-1295.

- Shear, M. K., Simon, N., Wall, M., Zisook, S., Neimeyer, R., Duan, N., . . . Keshaviah, A. (2011). Complicated grief and related bereavement issues for DSM-5. *Depression and Anxiety, 28*(2), 103-117. doi:10.1002/da.20780
- Smid, G. E., Kleber, R. J., de la Rie, S. M., Bos, J. B. A., Gersons, B. P. R., & Boelen, P. A. (2015). Brief eclectic psychotherapy for traumatic grief (BEP-TG): Toward integrated treatment of symptoms related to traumatic loss. *European Journal of Psychotraumatology, 6*(0), 1-11.
- Sprang, G. (2001). The use of eye movement desensitization and reprocessing (EMDR) in the treatment of traumatic stress and complicated mourning: Psychological and behavioral outcomes. *Research on Social Work Practice, 11*(3), 300-320. doi:10.1177/104973150101100302
- Stroebe, M., & Schut, H. (1999). The dual process model of coping with bereavement: Rationale and description. *Death Studies, 23*(3), 197-224.
- Stroud, C. B. (2018). The stress sensitization model. In K. Harkness, & E. P. Hayden (Eds.), *The oxford handbook of stress and mental health*. doi:10.1093/oxfordhb/9780190681777.013.16
- van der Velden, P. G., van der Meulen, E., Lenferink, L. I. M., & Ijzermans, J. C. (2018). Media experiences and associations with mental health among the bereaved of the MH17-disaster: A latent profile analysis. *Scandinavian Journal of Psychology, 59*(3), 281-288.
- WHO (2018). *ICD-11 beta draft. prolonged grief disorder criteria*. Retrieved from <https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/1183832314>.

Zisook, S., Shuchter, S. R., Pedrelli, P., Sable, J., & Deaciuc, S. C. (2001). Bupropion sustained release for bereavement: Results of an open trial. *The Journal of Clinical Psychiatry*, 62(4), 227-30.

In press

Supplemental Material Table 1
Fit indices of the cross-lagged models

PCBD and PTSD							
	χ^2	df	SABIC	AIC	RMSEA	CFI	TLI
Model 1	39.20	12	6241.46	6242.07	0.115	0.965	0.922
Model 2	41.03	16	6235.37	6235.90	0.095	0.968	0.946
Model 3	47.95	20	6234.36	6234.82	0.090	0.964	0.952
Model 4	100.44	23	6280.91	6281.31	0.140	0.901	0.884
PCBD and depression							
	χ^2	df	SABIC	AIC	RMSEA	CFI	TLI
Model 1	30.03	12	5634.64	5635.25	0.093	0.972	0.937
Model 2	33.25	16	5629.94	5630.47	0.079	0.973	0.955
Model 3	40.91	20	5629.68	5630.14	0.078	0.968	0.956
Model 4	87.76	23	5670.59	5670.99	0.128	0.899	0.882
PCBD, PTSD, and depression							
	χ^2	df	SABIC	AIC	RMSEA	CFI	TLI
Model 1	64.86	27	8461.81	8463.01	0.090	0.969	0.928
Model 2	71.63	33	8456.69	8457.78	0.082	0.968	0.940
Model 3	83.40	45	8444.69	8445.55	0.070	0.969	0.956
Model 4	175.56	54	8519.03	8519.71	0.114	0.901	0.884

Note. Model 1 includes autoregressive and cross-lagged paths and allows symptoms assessed at the same time point to covary. Model 2 resembles Model 1, but now we constrained the autoregressive effects to be equal for each distress measure separately. Model 3 resembles Model 2, but now we also constrained the cross-lagged effects to be equal across the waves. Model 4 resembles Model 3, but now we also constrained the associations between the distress measures at the same wave to be equal across the waves. The final models are shown in bold. AIC = Akaike information criterion; CFI = comparative fit index; df = degrees of freedom; PCBD = persistent complex bereavement disorder; PTSD = posttraumatic stress disorder; RMSEA = root mean square error of approximation; SA-BIC = sample-size adjusted Bayesian information criterion; TLI = Tucker-Lewis index.

Supplemental Material Table 2

Unstandardized autoregressive and cross-lagged effects and correlations for Model 3 (N = 172)

	B (SE)
PCBD and PTSD	
Constrained autoregressive effects	
PCBD	0.72*** (0.07)
PTSD	0.65*** (0.06)
Constrained cross-lagged effects	
PCBD on PTSD	0.08 (0.06)
PTSD on PCBD	0.20** (0.08)
Correlations	
PCBD W1 with PTSD W1	0.82 *** (0.03)
PCBD W2 with PTSD W2	0.83 *** (0.03)
PCBD W3 with PTSD W3	0.67*** (0.06)
PCBD W4 with PTSD W4	0.59*** (0.07)
PCBD and depression	
Constrained autoregressive effects	
PCBD	0.79*** (0.05)
Depression	0.66*** (0.05)
Constrained cross-lagged effects	
PCBD on depression	0.12 (0.12)
Depression on PCBD	0.05** (0.02)
Correlations	
PCBD W1 with depression W1	0.66*** (0.04)
PCBD W2 with depression W2	0.61*** (0.06)
PCBD W3 with depression W3	0.59*** (0.07)
PCBD W4 with depression W4	0.28** (0.10)
PCBD, PTSD, and depression	
Constrained autoregressive effects	
PCBD	0.72*** (0.07)
PTSD	0.56*** (0.07)
Depression	0.60*** (0.05)
Constrained cross-lagged effects	
PCBD on PTSD	0.07 (0.07)
PCBD on depression	0.08 (0.14)
PTSD on PCBD	0.18* (0.07)
PTSD on depression	0.43** (0.15)
Depression on PCBD	0.01 (0.03)
Depression on PTSD	0.06* (0.02)
Correlations	
PCBD W1 with PTSD W1	0.82*** (0.03)
PCBD W2 with PTSD W2	0.82*** (0.04)
PCBD W3 with PTSD W3	0.67*** (0.06)
PCBD W4 with PTSD W4	0.61*** (0.07)
PCBD W1 with depression W1	0.66*** (0.04)
PCBD W2 with depression W2	0.62*** (0.06)
PCBD W3 with depression W3	0.58*** (0.07)
PCBD W4 with depression W4	0.28** (0.10)

Depression W1 with PTSD W1	0.79*** (0.03)
Depression W2 with PTSD W2	0.67*** (0.06)
Depression W3 with PTSD W3	0.55*** (0.08)
Depression W4 with PTSD W4	0.44*** (0.08)

Note. PCBD = persistent complex bereavement disorder; PTSD = posttraumatic stress disorder. * $p < .05$ ** $p < .01$ *** $p < .001$

in press